Contents

Volume 114 1992

Couly, G. F., Coltey, P. M. and le Douarin, N. M.
The developmental fate of the cephalic mesoderm in quail-chick chimeras 1-15

Lafont, F., Rouget, M., Triller, A., Prochiantz, A. and Rousselet, A.
In vitro control of neuronal polarity by glycosaminoglycans 17-29

Hopwood, N. D., Pluck, A., Gurdon, J. B. and Dilworth, S. M.
Expression of XMyoD protein in early *Xenopus laevis* embryos 31-38

Sjöberg, M., Vennström, B. and Forrest, D.
Thyroid hormone receptors in chick retinal development: differential expression of mRNAs for α and N-terminal variant β receptors 39-47

Panzer, S., Weigel, D. and Beckendorf, S. K.
Organogenesis in *Drosophila melanogaster*: embryonic salivary gland determination is controlled by homeotic and dorsoventral patterning genes 49-57

Wood, T. L., Streck, R. D. and Pintar, J. E.
Expression of the IGFBP-2 gene in post-implantation rat embryos 59-66

Del Buono, R., Fleming, K. A., Morey, A. L., Hall, P. A. and Wright, N. A.
A nude mouse xenograft model of fetal intestine development and differentiation 67-23

Johnson, J. E., Zimmerman, K., Saito, T. and Anderson, D. J.
Induction and repression of mammalian *achaete-scute* homologue (MASH) gene expression during neuronal differentiation of P19 embryonal carcinoma cells 75-87

Gönczy, P., Viswanathan, S. and DiNardo, S.
Probing spermatogenesis in *Drosophila* with P-element enhancer detectors 89-98

Capovilla, M., Eldon, E. D. and Pirrotta, V.
The *giant* gene of *Drosophila* encodes a b-ZIP DNA-binding protein that regulates the expression of other segmentation gap genes 99-112

Raz, E. and Shilo, B.-Z.
Dissection of the *faint little ball (flb)* phenotype: determination of the development of the *Drosophila* central nervous system by early interactions in the ectoderm 113-123

Quéva, C., Ness, S. A., Grässer, F. A., Graf, T., Vandenbunder, B. and Stéhelin, D.
Expression patterns of *c-myb* and of *v-myb* induced myeloid-1 (*mim-1*) gene during the development of the chick embryo 125-133

Delarue, M., Sanchez, S., Johnson, K. E., Darribère, T. and Boucaut, J.-C.
A fate map of superficial and deep circumblastoporal cells in the early gastrula of *Pleurodeles waltl* 135-146

Lepage, T., Ghiglione, C. and Gache, C.
Spatial and temporal expression pattern during sea urchin embryogenesis of a gene coding for a protease homologous to the human protein BMP-1 and to the product of the *Drosophila* dorsal-ventral patterning gene *tolloid* 147-164

Hime, G. and Saint, R.
Zygotic expression of the *pebble* locus is required for cytokinesis during the postblastoderm mitoses of *Drosophila* 165-171

Goncharova, E. J., Kam, Z. and Geiger, B.
The involvement of adherens junction components in myofibrillogenesis in cultured cardiac myocytes 173-183

Alexiou, M. and Leese, H. J.
Purine utilisation, de novo synthesis and degradation in mouse preimplantation embryos 185-192

Nisson, P. E., Gaudette, M. F., Brandhorst, B. P. and Crain, W. R.
Mutually exclusive expression of the *Strongylocentrotus purpuratus* Spec1 gene and its *Lytechinus pictus* homologue in cells of hybrid embryos 193-201

Yancey, S. B., Biswal, S. and Revel, J.-P.
Spatial and temporal patterns of distribution of the gap junction protein connexin43 during mouse gastrulation and organogenesis 203-212

Nagel, M.-D. and Nagel, J.
Erythroid colony formation by fetal rat liver and spleen cells in vitro: inhibition by a low relative molecular mass component of fetal spleen 213-219

Contents

Macdonald, P. M.
The *Drosophila pumilio* gene: an unusually long transcription unit and an unusual protein 221-232

Peters, K. G., Werner, S., Chen, G. and Williams, L. T.
Two FGF receptor genes are differentially expressed in epithelial and mesenchymal tissues during limb formation and organogenesis in the mouse 233-243

Ohneda, O., Yanai, N. and Obinata, M.
Combined action of c-Kit and erythropoietin on erythroid progenitor cells 245-252

Lorenz, H. P., Longaker, M. T., Perkocha, L. A., Jennings, R. W., Harrison, M. R. and Adzick, N. S.
Scarless wound repair: a human fetal skin model 253-259

Kimelman, D. and Maas, A.
Induction of dorsal and ventral mesoderm by ectopically expressed *Xenopus* basic fibroblast growth factor 261-269

Gray, G. E. and Sanes, J. R.
Lineage of radial glia in the chicken optic tectum 271-283

ESSAY IN DEVELOPMENT
Slack, J. M. W. and Tannahill, D.
Mechanism of anteroposterior axis specificiation in vertebrates: lessons from the amphibians 285-302

CONTRIBUTED PAPERS
Wang, R., Clark, R. and Bautch, V. L.
Embryonic stem cell-derived cystic embryoid bodies form vascular channels: an in vitro model of blood vessel development 303-316

Bossinger, O. and Schierenberg, E.
Transfer and tissue-specific accumulation of cytoplasmic components in embryos of *Caenorhabditis elegans* and *Rhabditis dolichura*: in vivo analysis with a low-cost signal enhancement device 317-330

Watanabe, M., Kinutani, M., Naito, M., Ochi, O. and Takashima, Y.
Distribution analysis of transferred donor cells in avian blastodermal chimeras 331-338

Ordahl, C. P. and le Douarin, N. M.
Two myogenic lineages in the developing somite 339-353

Campos, A. R., Fischbach, K.-F. and Steller, H.
Survival of photoreceptor neurons in the compound eye of *Drosophila* depends on connections with the optic ganglia 355-366

Kitamura, K., Takiguchi-Hayashi, K., Sezaki, M., Yamamoto, H. and Takeuchi, T.
Avian neural crest cells express a melanogenic trait during early migration from the neural tube: observations with the new monoclonal antibody, "MEBL-1" 367-378

Carette, M. J. M. and Ferguson, M. W. J.
The fate of medial edge epithelial cells during palatal fusion in vitro: an analysis by DiI labelling and confocal microscopy 379-388

Kam, E. and Hodgins, M. B.
Communication compartments in hair follicles and their implication in differentiative control 389-393

Clerk, A., Strong, P. N. and Sewry, C. A.
Characterisation of dystrophin during development of human skeletal muscle 395-402

Selleck, M. A. J. and Stern, C. D.
Commitment of mesoderm cells in Hensen's node of the chick embryo to notochord and somite 403-415

Powell, B., Crocker, L. and Rogers, G.
Hair follicle differentiation: expression, structure and evolutionary conservation of the hair type II keratin intermediate filament gene family 417-433

Hirobe, T.
Basic fibroblast growth factor stimulates the sustained proliferation of mouse epidermal melanoblasts in a serum-free medium in the presence of dibutyryl cyclic AMP and keratinocytes 435-445

Behrendtsen, O., Alexander, C. M. and Werb, Z.
Metalloproteinases mediate extracellular matrix degradation by cells from mouse blastocyst outgrowths 447-456

Gagnon, M. L., Angerer, L. M. and Angerer, R. C.
Posttranscriptional regulation of ectoderm-specific gene expression in early sea urchin embryos 457-467

Lazzaro, D., De Simone, V., De Magistris, L., Lehtonen, E. and Cortese, R.
LFB1 and LFB3 homeoproteins are sequentially expressed during kidney development 469-479

Frank, L. H., Cheung, H.-K. and Cohen, R. S.
Identification and characterization of *Drosophila* female germ line transcriptional control elements 481-491

Simon, J., Chiang, A. and Bender, W.
Ten different *Polycomb* group genes are required for spatial control of the *abdA* and *AdbB* homeotic products 493-505

Chang, W. S., Serikawa, K., Allen, K. and Bentley, D.
Disruption of pioneer growth cone guidance in vivo by removal of glycosyl-phosphatidylinositol-anchored cell surface proteins 507-519

Breier, G., Albrecht, U., Sterrer, S. and Risau, W.
Expression of vascular endothelial growth factor during embryonic angiogenesis and endothelial cell differentitation 521-532

Müller, A. H. J., Angres, B. and Hausen, P.
U-cadherin in *Xenopus* oogenesis and oocyte maturation 533-543

Simon, A. M., Hoppe, P. and Burden, S. J.
Spatial restriction of AChR gene expression to subsynaptic nuclei 545-553

Ueno, K., Hui, C.-C., Fukuta, M. and Suzuki, Y.
Molecular analysis of the deletion mutants in the E homeotic complex of the silkworm *Bombyx mori* 555-563

Herzlinger, D., Koseki, C., Mikawa, T. and Al-Awqati, Q.
Metanephric mesenchyme contains multipotent stem cells whose fate is restricted after induction 565-572

Ihara, S. and Motobayashi, Y.
Wound closure in foetal rat skin 573-582

Ferguson, E. L. and Anderson, K. V.
Localized enhancement and repression of the activity of the TGF-β family member, *decapentaplegic*, is necessary for dorsal-ventral pattern formation in the *Drosophila* embryo 583-597

Bowman, J. L., Sakai, H., Jack, T., Weigel, D., Mayer, U. and Meyerowitz, E. M.
SUPERMAN, a regulator of floral homeotic genes in *Arabidopsis* 599-615

Grandin, N. and Charbonneau, M.
Intracellular free Ca^{2+} changes during physiological polyspermy in amphibian eggs 617-624

Taylor, B. J. and Truman, J. W.
Commitment of abdominal neuroblasts in *Drosophila* to a male or female fate is dependent on genes of the sex-determining hierarchy 625-642

Püschel, A. W., Gruss, P. and Westerfield, M.
Sequence and expression pattern of *pax-6* are highly conserved between zebrafish and mice 643-651

Cheung, H.-K., Serano, T. L. and Cohen, R. S.
Evidence for a highly selective RNA transport system and its role in establishing the dorsoventral axis of the *Drosophila* egg 653-661

Guthrie, S. and Lumsden, A.
Motor neuron pathfinding following rhombomere reversals in the chick embryo hindbrain 663-673

Spanakis, E., Lamina, P. and Bennett, D. C.
Effects of the developmental colour mutations silver and recessive spotting on proliferation of diploid and immortal mouse melanocytes in culture 675-680

Moffat, K. G., Gould, J. H., Smith, H. K. and O'Kane, C. J.
Inducible cell ablation in *Drosophila* by cold-sensitive ricin A chain 681-687

Rohrer, H.
Cholinergic neuronal differentiation factors: evidence for the presence of both CNTF-like and non-CNTF-like factors in developing rat footpad 689-698

Schroeder, M. M. and Gard, D. L.
Organization and regulation of cortical microtubules during the first cell cycle of *Xenopus* eggs 699-709

Isaacs, H. V., Tannahill, D. and Slack, J. M. W.
Expression of a novel FGF in the *Xenopus* embryo. A new candidate inducing factor for mesoderm formation and anteroposterior specification 711-720

di Clemente, N., Ghaffari, S., Pepinsky, R. B., Pieau, C., Josso, N., Cate, R. L. and Vigier, B.
A quantitative and interspecific test for biological activity of anti-Müllerian hormone: the fetal ovary aromatase assay 721-727

Storey, K. G., Crossley, J. M., De Robertis, E. M., Norris, W. E. and Stern, C. D.
Neural induction and regionalisation in the chick embryo 729-741

Guillemot, F. and Cepko, C. L.
Retinal fate and ganglion cell differentiation are potentiated by acidic FGF in an *in vitro* assay of early retinal development 743-754

Niswander, L. and Martin, G. R.
Fgf-4 expression during gastrulation, myogenesis, limb and tooth development in the mouse 755-768

Reynolds, S. D., Angerer, L. M., Palis, J., Nasir, A. and Angerer, R. C.
Early mRNAs, spatially restricted along the animal-vegetal axis of sea urchin embryos, include one encoding a protein related to tolloid and BMP-1 769-786

Bellen, H. J., D'Evelyn, D., Harvey, M. and Elledge, S. J.
Isolation of temperature-sensitive diphtheria toxins in yeast and their effects on *Drosophila* cells 787-796

Sanyal, S., Hawkins, R. K., Jansen, H. G. and Zeilmaker, G. H.
Compensatory synaptic growth in the rod terminals as a sequel to partial photoreceptor cell loss in the retina of chimaeric mice 797-803

Rowe, A., Richman, J. M. and Brickell, P. M.
Development of the spatial pattern of retinoic acid receptor-β transcripts in embryonic chick facial primordia 805-813

Chang, S., Fan, J. and Nayak, J.
Pathfinding by cranial nerve VII (facial) motorneurons in the chick hindbrain 815-823

Pike, S. H., Melancon, E. F. and Eisen, J. S.
Pathfinding by zebrafish motoneurons in the absence of normal pioneer axons 825-831

Miettinen, P. J. and Heikinheimo, K.
Transforming growth factor-alpha (TGF-α) and insulin gene expression in human fetal pancreas 833-840

Sundin, O. and Eichele, G.
An early marker of axial pattern in the chick embryo and its respecification by retinoic acid 841-852

Miner, J. H., Miller, J. B. and Wold, B. J.
Skeletal muscle phenotypes initiated by ectopic MyoD in transgenic mouse heart 853-860

Koshimizu, U., Watanabe, D., Tajima, Y. and Nishimune, Y.
Effects of W (c-*kit*) gene mutation on gametogenesis in male mice: agametic tubular segments in W^f/W^f testes 861-867

García-Aragón, J., Lobie, P. E., Muscat, G. E. O., Gobius, K. S., Norstedt, G. and Waters, M. J.
Prenatal expression of the growth hormone (GH) receptor/binding protein in the rat: a role for GH in embryonic and fetal development? 869-876

Castelli-Gair, J., Müller, J. and Bienz, M.
Function of an *Ultrabithorax* minigene in imaginal cells 877-886

Jahoda, C. A. B., Horne, K. A., Mauger, A., Bard, S. and Sengel, P.
Cellular and extracellular involvement in the regeneration of the rat lower vibrissa follicle 887-897

Watanabe, T. and Raff, M. C.
Diffusible rod-promoting signals in the developing rat retina 899-906

Thorp, B. H., Anderson, I. and Jakowlew, S. B.
Transforming Growth Factor-β1, -β2 and -β3 in cartilage and bone cells during endochondral ossification in the chick 907-911

Braisted, J. E. and Raymond, P. A.
Regeneration of dopaminergic neurons in goldfish retina 913-919

Pearce, J. J. H., Singh, P. B. and Gaunt, S. J.
The mouse has a *Polycomb*-like chromobox gene 921-929

Henry, J. J., Klueg, K. M. and Raff, R. A.
Evolutionary dissociation between cleavage, cell lineage and embryonic axes in sea urchin embryos 931-938

Skeath, J. B. and Carroll, S. B.
Regulation of proneural gene expression and cell fate during neuroblast segregation in the *Drosophila* embryo 939-946

Altshuler, D. and Cepko, C.
A temporally regulated, diffusible activity is required for rod photoreceptor development in vitro 947-957

Izpisúa-Belmonte, J.-C., Ede, D. A., Tickle, C. and Duboule, D.
The mis-expression of posterior *Hox-4* genes in *talpid* (ta^3) mutant wings correlates with the absence of anteroposterior polarity 959-963

Matzner, U., von Figura, K. and Pohlmann, R.
Expression of the two mannose 6-phosphate receptors is spatially and temporally different during mouse embryogenesis 965-972

Roark, E. F., Paradies, N. E., Lagunowich, L. A. and Grunwald, G. B.
Evidence for endogenous proteases, mRNA level and insulin as multiple mechanisms of N-cadherin down-regulation during retinal development 973-984

Hotary, K. B. and Robinson, K. R.
Evidence of a role for endogenous electrical fields in chick embryo development 985-996

Hagemann, L. J. and First, N. L.
Embryonic cytoplasmic extracts rescue murine androgenones to the blastocyst stage 997-1001

Arora, K. and Nüsslein-Volhard, C.
Altered mitotic domains reveal fate map changes in *Drosophila* embryos mutant for zygotic dorsoventral patterning genes 1003-1024

Index of Authors and Titles 1025-1028

Subject Index 1029-1041